

Lucidota eucera E. Olivier

Lucidota eucera E. Olivier, 1907. In Wytsman's Gen. Ins. 53:20.

Three males of this species were received from Peru; the species was described from French Guiana. It has an unusual development of the genital segment (fig. 3) and a relatively large aedeagus (fig. 4).



BOOK REVIEW

THE GROUND-BEETLES (CARABIDAE, EXCL. CICINDELINAE) OF CANADA AND ALASKA, PART 3. By Carl H. Lindroth. Opuscula Entomologica Suppl. 24:201-408, figs. 102-207, 1963. (Price, kr. 35:-; \$6.85.)

This volume continues the monograph of the Canadian and Alaskan Carabidae started in 1961 (see Coleopt. Bull. 16:106). The first part, the introduction, will be published after the text has been completed.

The importance of part 3 is apparent to any student of Coleoptera when it is pointed out that the work is concerned with the genus *Bembidion*. The 161 species treated include 16 new species, bringing the total for the genus in North America to about 411 species. Most of the species occur north of Washington, D. C., to about the tree line. Thus a substantial portion of the range of the genus is covered in this work. The care in which it is written and the detail of the study make it the standard work at the present time and it will replace all previous work for years to come.

It might well be added that any Coleopterist or entomology library that does not buy this series as issued will regret it when it is no longer available at the publisher's price.—ROSS H. ARNETT, JR.

LITERATURE NOTICE

CATALOGUE OF THE CANTHONINI (COL. SCARAB.) INHABITING THE WESTERN HEMISPHERE. By M. A. Vulcano and F. S. Pereira C.M.F. Ent. Arb. Mus. Frey 15(2):570-685. 1964.—This is a synonymical list with literature citations to every mention of each genus and species. Geographical distributions by countries, a bibliography of articles not mentioned by Blackwelder, and an index are also given.